

For each $\alpha > 2$ there is an infinite binary word with
critical exponent α

James D. Currie* & Narad Rampersad†
Department of Mathematics and Statistics
University of Winnipeg
Winnipeg, Manitoba R3B 2E9
CANADA
e-mail: currie@uwinnipeg.ca,
n.rampersad@uwinnipeg.ca

February 27, 2008

Abstract

For each $\alpha > 2$ there is a binary word with critical exponent α .

Keywords: Combinatorics on words, repetitions, critical exponent

Mathematics Subject Classifications: 68R15

1 Introduction

If α is a rational number, a word w is an α *power* if there exist words x and x' and a positive integer n , with x' a prefix of x , such that $w = x^n x'$ and $\alpha = n + |x'|/|x|$. We refer to $|x|$ as a *period* of w . A word is α *power-free* if none of its subwords is a β power with $\beta \geq \alpha$; otherwise, we say the word *contains an α power*.

The *critical exponent* of an infinite word $\mathbf{w}$ is defined as

$$\sup\{\alpha \in \mathbb{Q} \mid \mathbf{w} \text{ contains an } \alpha\text{-power}\}.$$

Critical exponents of certain classes of infinite words, such as Sturmian words [8, 10] and words generated by iterated morphisms [5, 6], have received particular attention.

Krieger and Shallit [7] proved that for every real number $\alpha > 1$, there is an infinite word with critical exponent α . As α tends to 1, the number of letters required to construct such words

*The author's research was supported by an NSERC operating grant.

†The author is supported by an NSERC Post-doctoral Fellowship.

tends to infinity. However, for $\alpha > 7/3$, Shur [9] gave a construction over a binary alphabet. For $\alpha > 2$, Krieger and Shallit gave a construction over a four-letter alphabet and left it as an open problem to determine if for every real number $\alpha \in (2, 7/3]$, there is an infinite binary word with critical exponent α . Currie, Rampersad, and Shallit [3] gave examples of such words for a dense subset of real numbers α in the interval $(2, 7/3]$. In this note we resolve the question completely by demonstrating that for every real number $\alpha > 2$, there is an infinite binary word with critical exponent α .

2 Properties of the Thue-Morse morphism

In this section we present some useful properties of the *Thue-Morse morphism*; i.e., the morphism μ defined by $\mu(0) = 01$ and $\mu(1) = 10$.

Lemma 1. *Let s be a positive integer. Let z be a subword of $\mu^s(01)$ with $|z| \geq 2^s$. Then z does not have period 2^s .*

Proof. Write $\mu^s(0) = a_1 a_2 \dots a_{2^n}$, $\mu^s(1) = b_1 b_2 \dots b_{2^n}$. One checks by induction that $a_i = 1 - b_i$ for $1 \leq i \leq 2^n$, and the result follows. $\square$

Brandenburg [1] proved the following useful theorem, which was independently rediscovered by Shur [9].

Theorem 2 (Brandenburg; Shur). *Let w be a binary word and let $\alpha > 2$ be a real number. Then w is α power-free if and only if $\mu(w)$ is α power-free.*

The following sharper version of one direction of this theorem (implicit in [4]) is also useful.

Theorem 3. *Suppose $\mu(w)$ contains a subword u of period p , with $|u|/p > 2$. Then w contains a subword v of length $\lceil |u|/2 \rceil$ and period $p/2$.*

We will also have call to use the deletion operator δ which removes the first (left-most) letter of a word. For example, $\delta(12345) = 2345$.

3 A binary word with critical exponent α

We denote by $\mathcal{L}$ the set of factors (subwords) of words of $\mu(\{0, 1\}^*)$.

Lemma 4. *Let $00v \in \mathcal{L}$, and suppose that $00v$ is α power-free, some fixed $\alpha > 2$. Let $r = \lceil \alpha \rceil$. Suppose that $0^r v = xuy$ where u is an α power. Then $x = \epsilon$ and $u = 0^r$.*

Proof. Suppose that u has period p . Since $00v$ is α power-free, we can write $u = 0^s v'$, $x = 0^{r-s}$ for some integer s , $3 \leq s \leq r$, some prefix v' of v . If 0^p is not a prefix of u then the prefix of u of length p contains the subword 0001 . Since $\alpha > 2$, this means that 0001 is a subword of u at least twice, so that 0001 is a subword of $00v$. This is impossible, since $00v \in \mathcal{L}$.

Therefore, 0^p is a prefix of u , and u has the form 0^t , some integer $t \geq \alpha$. This implies that u has 0^r as a prefix, so that $x = \epsilon$ and $u = 0^r$. $\square$

Lemma 5. *Let $\alpha > 2$ be given, and let $r = \lceil \alpha \rceil$. Let s, t be positive integers, such that $s \geq 3$ and there are words $x, y \in \{0, 1\}^*$ such that $\mu^s(0) = x00y$ with $|x| = t$. Suppose that $2 < r - t/2^s < \alpha$ and $00v \in \mathcal{L}$ is α power-free.*

1. *The word $\delta^t \mu^s(0^r w)$ has a prefix which is a β power, where $\beta = r - t/2^s$.*
2. *Suppose that $00v$ contains a β power of period p , some β and p . Then $\delta^t \mu^s(0^r w)$ contains a β power of period $2^s p$.*
3. *Word $\delta^t \mu^s(0^r w)$ is α power-free*

Proof. We start by observing that $\mu^s(0^r)$ has period 2^s . It follows that $\delta^t \mu^s(0^r)$ has period 2^s , length $r2^s - t$, and hence is a $(r2^s - t)/2^s = \beta$ power.

Now suppose u is a β power of period p in $00v$. Then $\mu(u)$ is a β power of period 2^s in $\mu^s(00v)$. However, $\mu^s(0^{r-1}v)$ is a suffix of $\delta^t \mu^s(0^r v)$, since $t < 2^s = |\mu^s(0)|$. Thus $\mu^s(u)$ is a β power of period 2^s in $\delta^t \mu^s(0^r v)$.

Next, note that $\mu^s(0^{r-1}v)$ does not contain any κ power, $\kappa \geq \alpha$. Otherwise, by Theorem 3 and induction, $0^{r-1}v$ contains a κ power. This is impossible by Lemma 4.

Suppose then that $\delta^t \mu^s(0^r v)$ contains a κ power $\hat{u}$ of period p , $\kappa \geq \alpha$. Using induction and Theorem 3, $0^r v$ contains a κ power u of period $p/2^s$. By Lemma 4, the only possibility is $u = 0^r$, and $p/2^s = 1$. Thus $p = 2^s$.

Since $00v \in \mathcal{L}$, the first letter of v is a 1. Since $\hat{u}$ has period 2^s , by Lemma 1 no subword of $\mu^s(01)$ of length greater than 2^s occurs in $\hat{u}$. We conclude that either $\hat{u}$ is a subword of $\delta^t \mu^s(0^r)$, or of $\mu^s(v)$, and hence of $\mu^s(0^{r-1}v)$. As this second case has been ruled out earlier, we conclude that $|\hat{u}| \leq |\delta^t \mu^s(0^r)| = r2^s - t$. This gives a contradiction: $\hat{u}$ is a κ power, yet $|\hat{u}|/p \leq (r2^s - t)/2^s = \beta < \alpha$. $\square$

By construction, $\delta^t \mu^s(0^r w)$ has the form $00\hat{v}$ where $00\hat{v} \in \mathcal{L}$.

We are now ready to prove our main theorem:

Theorem 6. *Let $\alpha > 2$ be a real number. There is a word over $\{0, 1\}$ with critical exponent α .*

Proof. Call a real number $\beta < \alpha$ *obtainable* if β can be written $\beta = r - t/2^s$, where r, s, t are positive integers, $s \geq 3$, and the word obtained by removing a prefix of length t from $\mu^s(0)$ begins with 00 . We note that $\mu^3(0) = 01101001$ and $\mu^3(1) = 10010110$ are of length 8, and both contain 00 as a subword; for a given $s \geq 3$ it follows that r and t can be chosen so that $\beta = r - t/2^s < \alpha$ and $|\alpha - \beta| \leq 7/2^s$; by choosing large enough s , an obtainable number β can be chosen arbitrarily close to α .

Let $\{\beta_i\}$ be a sequence of obtainable numbers converging to α . For each i write $\beta_i = r_i - t_i/2^{s_i}$, where r_i, s_i, t_i are positive integers, $s_i \geq 3$, and the word obtained by removing a prefix of length t_i from $\mu^{s_i}(0)$ begins with 00 . If $00w \in \mathcal{L}$, denote by $\phi_i(w)$ the word $\delta^{t_i} \mu^{s_i}(0^{r_i} w)$.

Consider the sequence of words

$$\begin{aligned}
w_1 &= \phi_1(\epsilon) \\
w_2 &= \phi_1(\phi_2(\epsilon)) \\
w_3 &= \phi_1(\phi_2(\phi_3(\epsilon))) \\
&\vdots \\
w_n &= \phi_1(\phi_2(\phi_3(\cdots(\phi_n(\epsilon))\cdots))) \\
&\vdots
\end{aligned}$$

By the third part of Lemma 5, if $00w \in \mathcal{L}$ is α power-free, then so is $\phi_i(w)$. Since 00ϵ is α power-free, each w_i is therefore α power-free.

By the first and second parts of Lemma 5, w_n contains β_i powers, $i = 1, 2, \dots, n$.

Note that ϵ is a prefix of $\phi_{n+1}(\epsilon)$, so that

$$w_n = \phi_1(\phi_2(\phi_3(\cdots(\phi_n(\epsilon))\cdots)))$$

is a prefix of

$$\phi_1(\phi_2(\phi_3(\cdots(\phi_n(\phi_{n+1}(\epsilon))\cdots))) = w_{n+1}.$$

We may therefore let $w = \lim_{n \rightarrow \infty} w_i$.

Since every prefix of w is α power-free, w is α power-free but contains β_i powers for each i . The critical exponent of w is therefore α . $\square$

The following question raised by Krieger and Shallit remains open: for $\alpha > 1$, if α powers are avoidable on a k -letter alphabet, does there exist an infinite word over k letters with critical exponent α ? In particular, for $\alpha > \text{RT}(k)$, where $\text{RT}(k)$ denotes the *repetition threshold* on k letters (see [2]), does there exist an infinite word over k letters with critical exponent α ? We believe that the answer is “yes”.

References

- [1] F.-J. Brandenburg, “Uniformly growing k -th power-free homomorphisms”, *Theoret. Comput. Sci.* **23** (1983), 69–82.
- [2] A. Carpi, “On Dejean’s conjecture over large alphabets”, *Theoret. Comput. Sci.* **385** (2007), 137–151.
- [3] J.D. Currie, N. Rampersad, J. Shallit, “Binary words containing infinitely many overlaps”, *Electron. J. Combin.* **13** (2006), #R82.
- [4] J. Karhumäki, J. Shallit, “Polynomial versus exponential growth in repetition-free binary words”, *J. Combin. Theory Ser. A* **104** (2004), 335–347.
- [5] D. Krieger, “On critical exponents in fixed points of binary k -uniform morphisms”. In *Proc. STACS 2006*, LNCS 3884, Springer-Verlag, 2006, pp. 104–114.

- [6] D. Krieger, “On critical exponents in fixed points of non-erasing morphisms”, *Theoret. Comput. Sci.* **376** (2007), 70-88.
- [7] D. Krieger, J. Shallit, “Every real number greater than 1 is a critical exponent”, *Theoret. Comput. Sci.* **381** (2007), 177-182.
- [8] F. Mignosi, G. Pirillo, “Repetitions in the Fibonacci infinite word”, *RAIRO Inform. Theor. Appl.* **26** (1992), 199-204.
- [9] A. M. Shur, “The structure of the set of cube-free $\mathbb{Z}$ -words in a two-letter alphabet” (Russian), *Izv. Ross. Akad. Nauk Ser. Mat.* **64** (2000), 201–224. English translation in *Izv. Math.* **64** (2000), 847–871.
- [10] D. Vandeth, “Sturmian words and words with a critical exponent”, *Theoret. Comput. Sci.* **242** (2000), 283-300.